

Remark on a conjecture of Mukai

Arnaud BEAUVILLE

Introduction

The conjecture mentioned in the title appears actually as a question in [M] (Problem 4.11):

Conjecture .— *Let C be a general curve, and E a stable rank 2 vector bundle on C with $\det E = K_C$. The multiplication map $\mu_E : \mathbf{S}^2 H^0(C, E) \rightarrow H^0(C, \mathbf{S}^2 E)$ is injective.*

Let $\mathcal{M}_K$ be the moduli space of stable rank 2 vector bundles E on C with $\det E = K_C$, and let $\mathcal{M}_K^n$ be the subvariety of $\mathcal{M}_K$ parametrizing bundles with $h^0(E) = n$. As explained in *loc. cit.*, the conjecture implies that $\mathcal{M}_K^n$ is smooth, of codimension $\frac{1}{2}n(n+1)$; this would give an analogue of the Brill-Noether theory for rank 2 vector bundles with canonical determinant.

In this note we prove the conjecture in a very particular case:

Proposition 1 .— *The conjecture holds if $h^0(E) \leq 6$.*

As a corollary we obtain that $\mathcal{M}_K^n$ is smooth of codimension $\frac{1}{2}n(n+1)$ in $\mathcal{M}$ for $n \leq 6$. Another consequence is that Mukai's conjecture holds for $g \leq 9$ (see § 2).

The idea of the proof is the following. Put $n = h^0(E)$; assume for simplicity that E is generated by its global sections, and that the map μ_E annihilates a non-degenerate (symmetric) tensor. This means that the image of the map $\mathbf{P}_C(E) \rightarrow \mathbf{P}^{n-1}$ defined by the tautological line bundle is contained in a smooth quadric; equivalently, the map from C into the Grassmann variety $\mathbf{G}(2, n)$ associated to E factors through the orthogonal Grassmannian $\mathbf{GO}(2, n)$. Now for $n \leq 6$, the restriction of the Plücker line bundle $\mathcal{O}_{\mathbf{G}}(1)$ to $\mathbf{GO}(2, n)$ is the sum of two (possibly equal) globally generated line bundles; pulling back to C we obtain a decomposition of $\det E = K_C$ which turns out to contradict Brill-Noether theory for a general curve.

The last part of the argument breaks down for $n \geq 7$, since then the restriction map $\mathrm{Pic}(\mathbf{G}(2, n)) \rightarrow \mathrm{Pic}(\mathbf{GO}(2, n))$ is an isomorphism. In fact we will show that the result cannot be improved without strengthening the hypotheses that we use on C and E .

1. Proof of the main result

We will say that a curve C is *Brill-Noether general* if for any line bundle L

on C , the multiplication map

$$H^0(C, L) \otimes H^0(C, K_C \otimes L^{-1}) \longrightarrow H^0(C, K_C)$$

is injective. A general curve of given genus is Brill-Noether general; if S is a K3 surface with $\text{Pic}(S) = \mathbf{Z}[H]$, a general element of the linear system $|H|$ is Brill-Noether general [L].

We will use this property in the following way:

Lemma 1.— *Let C be a Brill-Noether general curve and L, L' two line bundles on C such that $h^0(K_C \otimes (L \otimes L')^{-1}) \geq 1$. Then :*

- a) *The multiplication map $H^0(C, L) \otimes H^0(C, L') \rightarrow H^0(C, L \otimes L')$ is injective;*
- b) *If $L' = L$, we have $h^0(L) \leq 1$.*

Proof : Choose a non-zero section $s \in H^0(C, (K_C \otimes (L \otimes L')^{-1}))$, and consider the commutative diagram

$$\begin{array}{ccc} H^0(L) \otimes H^0(L') & \longrightarrow & H^0(L \otimes L') \\ 1 \otimes s \downarrow & & \downarrow s \\ H^0(L) \otimes H^0(K_C \otimes L^{-1}) & \longrightarrow & H^0(K_C) . \end{array}$$

By our hypothesis on C the bottom horizontal map is injective; it follows that the top one is injective.

If $L' = L$, we get that the map $H^0(L)^{\otimes 2} \rightarrow H^0(L^{\otimes 2})$ is injective; if $H^0(L)$ contains two linearly independent elements s, t , the tensor $s \otimes t - t \otimes s$ is non-zero and belongs to the kernel of that map, a contradiction. ■

Let E be a vector bundle on a curve C . If the multiplication map μ_E is injective, the same property holds for all subbundles of E ; thus E satisfies

($\star$) *For every sub-line bundle $L \subset E$, the map $\mu_L : \mathbf{S}^2 H^0(C, L) \rightarrow H^0(C, L^2)$ is injective.*

By [T], this condition is automatically satisfied if C is general and any sub-line bundle of E has degree $\leq g + 1$. Thus Proposition 1 is a consequence of the following more precise result:

Proposition 2.— *Let C be a Brill-Noether general curve, and E a rank 2 vector bundle on C with $\det E = K_C$, satisfying condition ($\star$). Then any non-zero tensor $\tau \in \mathbf{S}^2 H^0(C, E)$ such that $\mu_E(\tau) = 0$ has rank > 6 .*

Proof : a) *The general set-up*

Let C be a curve, E a rank 2 vector bundle on C with $\det E = K_C$, τ an element of $\text{Ker } \mu_E$ of rank n . This means that we can find linearly independent

elements $s_1, \dots, s_n$ of $H^0(C, E)$ such that $\tau = s_1^2 + \dots + s_n^2$. Let $F \subset E$ be the image of the map $\mathcal{O}_C^n \rightarrow E$ defined by $s_1, \dots, s_n$. Then $s_1, \dots, s_n$ are sections of F which satisfy $s_1^2 + \dots + s_n^2 = 0$ in $H^0(C, \mathbf{S}^2 F)$. By property $(\star)$ F is a rank 2 subsheaf of E , with determinant $K_C(-A)$ for some effective divisor A .

Let $P := \mathbf{P}_C(F)$, and let $\mathcal{O}_P(1)$ be the tautological line bundle on P . Through the canonical isomorphisms

$$H^0(P, \mathcal{O}_P(1)) \xrightarrow{\sim} H^0(C, F) \quad H^0(P, \mathcal{O}_P(2)) \xrightarrow{\sim} H^0(C, \mathbf{S}^2 F)$$

the multiplication map μ_F is identified with $\mu_{\mathcal{O}_P(1)}$; thus we can view $s_1, \dots, s_n$ as global sections of $\mathcal{O}_P(1)$, which generate $\mathcal{O}_P(1)$ and satisfy $s_1^2 + \dots + s_n^2 = 0$. In other words, the image of the morphism $\varphi : P \rightarrow \mathbf{P}^{n-1}$ defined by $(s_1, \dots, s_n)$ is contained in the smooth quadric Q defined by $X_1^2 + \dots + X_n^2 = 0$.

Let γ be the map of C into the Grassmann variety $\mathbf{G}(2, n)$ associated to the surjective homomorphism $\mathcal{O}_C^n \rightarrow F$; by definition F is the pull back of the universal quotient bundle on $\mathbf{G}(2, n)$. The determinant of that bundle is the Plücker line bundle $\mathcal{O}_{\mathbf{G}}(1)$ on $\mathbf{G}(2, n)$, so we get $\gamma^* \mathcal{O}_{\mathbf{G}}(1) \cong K_C(-A)$.

For each $x \in C$, the point $\gamma(x) \in \mathbf{G}(2, n)$ corresponds to the line $\varphi(\mathbf{P}(F_x))$ in $\mathbf{P}^{n-1}$. So the fact that the image of φ is contained in Q means that γ factors through the orthogonal grassmannian $\mathbf{GO}(2, n)$ of lines contained in Q :

$$\gamma : C \xrightarrow{\gamma_{\mathcal{O}}} \mathbf{GO}(2, n) \hookrightarrow \mathbf{G}(2, n) .$$

b) *The cases $n = 4$ and $n = 5$*

Now we will assume $n \leq 6$ and that C is Brill-Noether general, and derive a contradiction. The case $n \leq 3$ is trivial ($\mathbf{GO}(2, n)$ is empty). Consider the case $n = 4$. Then $\mathbf{GO}(2, 4)$ parametrizes the lines in a smooth quadric in $\mathbf{P}^3$; it has two components, both isomorphic to $\mathbf{P}^1$. Inside $\mathbf{G}(2, 4)$, which is a quadric in $\mathbf{P}^5$, each of these components is a conic. Thus γ factors as

$$\gamma : C \xrightarrow{\gamma_{\mathcal{O}}} \mathbf{P}^1 \hookrightarrow \mathbf{G}(2, 4) ,$$

with $\mathcal{O}_{\mathbf{G}}(1)|_{\mathbf{P}^1} = \mathcal{O}_{\mathbf{P}^1}(2)$. Thus $L = \gamma_{\mathcal{O}}^* \mathcal{O}_{\mathbf{P}^1}(1)$ satisfies $L^2 = K_C(-A)$ and $h^0(L) \geq 2$, contradicting Lemma 1 b).

Suppose $n = 5$. Let V be a 4-dimensional vector space, with a non-degenerate alternate form ω and an orientation $\Lambda^4 V \xrightarrow{\sim} \mathbf{C}$. The orthogonal W of ω in $\Lambda^2 V$ is a 5-dimensional vector space with a non-degenerate quadratic form given by the wedge product. The isotropic 2-planes in W are of the form $\ell \wedge \ell^\perp$, for $\ell \in \mathbf{P}(V)$. Thus $\mathbf{GO}(2, 5) \subset \mathbf{G}(2, 5)$ is identified with $\mathbf{P}(V)$ embedded in $\mathbf{G}(2, W)$ by $\ell \mapsto \ell \wedge \ell^\perp$. The corresponding map from $\mathbf{P}(V)$ to $\mathbf{P}(\Lambda^2 W)$ is quadratic, so again $\mathcal{O}_{\mathbf{G}}(1)|_{\mathbf{P}(V)} = \mathcal{O}_{\mathbf{P}(V)}(2)$, and we conclude exactly as above.

c) *The case $n = 6$*

Let again V be a 4-dimensional vector space with an orientation. Then $\Lambda^2 V$ is a 6-dimensional quadratic vector space; the corresponding quadric in $\mathbf{P}(V)$ is the Grassmannian $\mathbf{G}(2, V)$. The lines contained in $\mathbf{G}(2, V) \subset \mathbf{P}(\Lambda^2 V)$ correspond to the 2-planes $\ell \wedge H \subset \Lambda^2 V$ where $\ell \subset H \subset V$, $\dim \ell = 1$, $\dim H = 3$; thus $\mathbf{GO}(2, 6) \subset \mathbf{G}(2, 6)$ is identified with the incidence variety $Z \subset \mathbf{P}(V) \times \mathbf{P}(V^*)$, embedded in $\mathbf{G}(2, V)$ by $(\ell, H) \mapsto \ell \wedge H$. Fixing ℓ or H one sees that the restriction of $\mathcal{O}_{\mathbf{G}}(1)$ to Z is the pull back of $\mathcal{O}_{\mathbf{P}}(1) \boxtimes \mathcal{O}_{\mathbf{P}}(1)$.

The map $\gamma_{\mathbf{O}} : C \rightarrow Z$ gives by projection two maps $u : C \rightarrow \mathbf{P}(V)$ and $u' : C \rightarrow \mathbf{P}(V^*)$. Put $L = u^* \mathcal{O}_{\mathbf{P}(V)}(1)$, $L' = u'^* \mathcal{O}_{\mathbf{P}(V^*)}(1)$. Then $L \otimes L' \cong K_C(-A)$, and we deduce from u and u' two homomorphisms

$$v : V^* \rightarrow H^0(C, L) \quad v' : V \rightarrow H^0(C, L') .$$

Consider the map

$$V \otimes V^* \xrightarrow{v \otimes v'} H^0(L) \otimes H^0(L') \longrightarrow H^0(L \otimes L') ;$$

the fact that $(u, u') : C \rightarrow \mathbf{P}(V) \times \mathbf{P}(V^*)$ factors through Z means that the identity element of $V \otimes V^* \cong \text{End}(V)$ goes to zero in $H^0(L \otimes L')$, and therefore already in $H^0(L) \otimes H^0(L')$ by Lemma 1 a).

Put $K = \text{Ker } u$, $K' = \text{Ker } u'$. The kernel of $v \otimes v'$ is $K \otimes V^* + V \otimes K'$; any element of this kernel has rank $\leq \dim K + \dim K'$. Since the identity tensor has rank 4, we get $\dim K + \dim K' \geq 4$.

Suppose $\dim K = \dim K' = 2$. Identifying again $V \otimes V^*$ to $\text{End}(V)$, we can write $\text{Id}_V = p + q$, with $\text{Im } p \subset K$ and $\text{Ker } q \supset K'^{\perp}$. This implies that p and q are orthogonal projectors, and therefore that $K' = K^{\perp}$. Then $\mathbf{P}(K) \times \mathbf{P}(K^{\perp})$ is contained in Z , and $\gamma_{\mathbf{O}}$ factors as

$$C \xrightarrow{(u, u')} \mathbf{P}(K) \times \mathbf{P}(K^{\perp}) \hookrightarrow \mathbf{GO}(2, 6) .$$

Let $(\ell, H) \in \mathbf{P}(K) \times \mathbf{P}(K^{\perp})$; then $\ell \subset K$ and $H \supset K$, so the line $\mathbf{P}(\ell \wedge H)$ in $\mathbf{P}(\Lambda^2 V) = \mathbf{P}^5$ contains the point $p := \mathbf{P}(\Lambda^2 K)$. In other words, the lines in Q parametrized by C all pass through p , hence are contained in the tangent hyperplane $T_p(Q)$. Therefore the scroll $\varphi(P)$ in $\mathbf{P}^5$ is contained in $T_p(Q)$; this is impossible because φ is defined by 6 linearly independent sections of $\mathcal{O}_{\mathbf{P}}(1)$.

Suppose now $\dim K = 3$, so that the image of C in $\mathbf{P}(V^*)$ is the point $\mathbf{P}(K^{\perp})$. Then $\gamma_{\mathbf{O}}$ factors as

$$C \xrightarrow{u} \mathbf{P}(K) \hookrightarrow \mathbf{GO}(2, 6) ;$$

the image of $\mathbf{P}(K)$ in $\mathbf{GO}(2, 6)$ is the family of lines contained in $\mathbf{P}(\Lambda^2 K) \subset \mathbf{P}(\Lambda^2 V) = \mathbf{P}^5$. Thus φ maps P to the projective plane $\mathbf{P}(\Lambda^2 K)$, which is again impossible. The same argument applies when $\dim K' = 3$. ■

2. Consequences and comments

The following lemma must be well-known:

Lemma 2.— *Let C be a Brill-Noether general curve, and E a semi-stable rank 2 vector bundle on C with $\det E = K_C$. Then $h^0(E) \leq \frac{g}{2} + 2$.*

Proof: Let L be a sub-line bundle of E of maximal degree. The exact sequence

$$0 \rightarrow L \rightarrow E \rightarrow K_C \otimes L^{-1} \rightarrow 0$$

gives $h^0(E) \leq h^0(L) + h^0(K_C \otimes L^{-1})$. Since $h^0(L)h^0(K_C \otimes L^{-1}) \leq g$ and $h^0(L) \leq h^0(K_C \otimes L^{-1})$ by semi-stability, the required inequality holds unless $h^0(L)$ is 0 or 1. In that case we have

$$h^0(L) + h^0(K_C \otimes L^{-1}) = 2h^0(L) + g - 1 - \deg L \leq g + 1 - \deg L$$

and $\deg L \geq \frac{g}{2} - 1$ by [N], hence $h^0(E) \leq \frac{g}{2} + 2$. ■

In particular $g \leq 9$ guarantees $h^0(E) \leq 6$, hence:

Corollary.— *Mukai's conjecture holds for $g \leq 9$.*

We will show that Proposition 2 cannot be improved without further hypotheses:

Proposition 3.— *For each integer $s \geq 5$, there exists a Brill-Noether general curve C of genus $g = 2s$ and a stable rank 2 vector bundle E on C with $\det E = K_C$, satisfying $(\star)$, such that $\text{Ker } \mu_E$ contains a tensor of rank 7.*

Proof: We choose a K3 surface S with $\text{Pic}(S) = \mathbf{Z}[H]$, $(H^2) = 4s - 2$, and take for C a general element of the linear system $|H|$. Then C has genus $2s$ and is Brill-Noether general; in particular it admits a finite set $\mathcal{P}$ of line bundles L with $h^0(L) = 2$, $\deg L = s + 1$. Choose one of these line bundles, say L ; the *Lazarsfeld bundle* E_L is defined by the exact sequence

$$0 \rightarrow E_L^* \rightarrow H^0(C, L) \otimes \mathcal{O}_S \rightarrow L \rightarrow 0.$$

It turns out that the restriction E of E_L to C does not depend on the choice of $L \in \mathcal{P}$; we have $\det E = K_C$, $h^0(E) = s + 2$ (see [V]), and for each $L \in \mathcal{P}$ an exact sequence

$$0 \rightarrow L \rightarrow E \rightarrow K_C \otimes L^{-1} \rightarrow 0.$$

Lemma 3.— *Let M be a sub-line bundle of E . Then either $M \in \mathcal{P}$, or $h^0(M) \leq 1$ and $\deg M \leq s$.*

Proof: Put $M' := K_C \otimes M^{-1}$; we may assume that the E/M is isomorphic to M' , so that

$$s + 2 = h^0(E) \leq h^0(M) + h^0(M') = 2h^0(M) + g - 1 - \deg M = 2h^0(M') + g - 1 - \deg M'.$$

As in the proof of Lemma 2 this implies that either $h^0(M)$ or $h^0(M')$ is ≤ 2 .

If $h^0(M) \leq 2$, we get $\deg M \leq s + 1$, hence either $M \in \mathcal{P}$, or $\deg M \leq s$ and $h^0(M) \leq 1$.

Assume $h^0(M') \leq 2$, so that $h^0(M) \geq s$. We get again $\deg M' \leq s + 1$, hence $\deg M \geq 3s - 3$. On the other hand the exact sequence preceding the Lemma shows that M injects into $K_C \otimes L^{-1}$ for each $L \in \mathcal{P}$; since $\text{Card}(\mathcal{P}) \geq 2$ the inclusion must be strict, so $\deg M < \deg K_C \otimes L^{-1} = 3s - 3$, a contradiction. ■

It follows that E is stable and satisfies condition $(\star)$. We observe that $\mathbf{S}^2 E$ is isomorphic to $K_C \otimes \mathcal{E}nd_0(E)$, where $\mathcal{E}nd_0(E)$ is the sheaf of trace 0 endomorphisms of E ; thus Serre duality and the stability of E imply $h^1(\mathbf{S}^2 E) = h^0(\mathcal{E}nd_0(E)) = 0$, hence $h^1(\mathbf{S}^2 E) = 3g - 3$ by Riemann-Roch.

In $\mathbf{S}^2 H^0(E)$ the locus of tensors of rank ≤ 7 has dimension $7(s - 1)$, while the kernel of μ_E has codimension $\leq 6s - 3$. Therefore the intersection of these subvarieties is not reduced to 0 – in fact it has dimension $\geq s - 4$. By Proposition 2 all the tensors in this intersection have rank 7. ■

REFERENCES

- [L] R. LAZARSELD: *Brill-Noether-Petri without degenerations*. J. Differential Geom. **23** (1986), no. 3, 299–307.
- [M] S. MUKAI: *Vector bundles and Brill-Noether theory*. Current topics in complex algebraic geometry, 145–158; Math. Sci. Res. Inst. Publ. **28**, Cambridge Univ. Press, 1995.
- [N] M. NAGATA: *On self-intersection number of a section on a ruled surface*. Nagoya Math. J. **37** (1970), 191–196.
- [T] M. TEIXIDOR: *Injectivity of the symmetric map for line bundles*. Manuscripta Math. **112** (2003), no. 4, 511–517.
- [V] C. VOISIN: *Sur l'application de Wahl des courbes satisfaisant la condition de Brill-Noether-Petri*. Acta Math. **168** (1992), no. 3-4, 249–272.

Arnaud BEAUVILLE
Institut Universitaire de France
&
Laboratoire J.-A. Dieudonné
UMR 6621 du CNRS
UNIVERSITÉ DE NICE
Parc Valrose
F-06108 NICE Cedex 02